

Bifurcation and multiplicity results for critical fractional p -Laplacian problems*

Kanishka Perera

Department of Mathematical Sciences
Florida Institute of Technology
Melbourne, FL 32901, USA

Marco Squassina[†]

Dipartimento di Informatica
Università degli Studi di Verona
37134 Verona, Italy

Yang Yang[‡]

School of Science
Jiangnan University
Wuxi, 214122, China

Abstract

We prove a bifurcation and multiplicity result for a critical fractional p -Laplacian problem that is the analog of the Brézis-Nirenberg problem for the nonlocal quasilinear case. This extends a result in the literature for the semilinear case $p = 2$ to all $p \in (1, \infty)$, in particular, it gives a new existence result. When $p \neq 2$, the nonlinear operator $(-\Delta)_p^s$, $s \in (0, 1)$ has no linear eigenspaces, so our extension is nontrivial and requires a new abstract critical point theorem that is not based on linear subspaces. We prove a new abstract result based on a pseudo-index related to the \mathbb{Z}_2 -cohomological index that is applicable here.

**MSC2010*: Primary 35R11, 35B33, Secondary 35B32, 58E05

Key Words and Phrases: fractional p -Laplacian, critical nonlinearity, bifurcation, multiplicity, existence, abstract critical point theory, \mathbb{Z}_2 -cohomological index, pseudo-index

[†]The second-named author was supported by 2009 MIUR project: “Variational and Topological Methods in the Study of Nonlinear Phenomena”.

[‡]This work was completed while the third-named author was visiting the Department of Mathematical Sciences at the Florida Institute of Technology, and she is grateful for the kind hospitality of the department. Project supported by NSFC-Tian Yuan Special Foundation (No. 11226116), Natural Science Foundation of Jiangsu Province of China for Young Scholars (No. BK2012109), and the China Scholarship Council (No. 201208320435).

1 Introduction and main results

For $p \in (1, \infty)$, $s \in (0, 1)$, and $N > sp$, the fractional p -Laplacian $(-\Delta)_p^s$ is the nonlinear nonlocal operator defined on smooth functions by

$$(-\Delta)_p^s u(x) = 2 \lim_{\varepsilon \searrow 0} \int_{\mathbb{R}^N \setminus B_\varepsilon(x)} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))}{|x - y|^{N+sp}} dy, \quad x \in \mathbb{R}^N.$$

This definition is consistent, up to a normalization constant depending on N and s , with the usual definition of the linear fractional Laplacian operator $(-\Delta)^s$ when $p = 2$. There is currently a rapidly growing literature on problems involving these nonlocal operators. In particular, fractional p -eigenvalue problems have been studied in Brasco and Parini [6], Brasco, Parini and Squassina [5], Franzina and Palatucci [20], Iannizzotto and Squassina [28] and in Lindgren and Lindqvist [31]. Regularity of solutions was obtained in Di Castro, Kuusi and Palatucci [14, 15], Iannizzotto, Mosconi and Squassina [26], Kuusi, Mingione and Sire [29] and Lindgren [30]. Existence via Morse theory was investigated in Iannizzotto et al. [27]. We refer to Caffarelli [9] for the motivations that have lead to their study. Let Ω be a bounded domain in \mathbb{R}^N with Lipschitz boundary. We consider the problem

$$\begin{cases} (-\Delta)_p^s u = \lambda |u|^{p-2} u + |u|^{p_s^*-2} u & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^N \setminus \Omega, \end{cases} \quad (1.1)$$

where $p_s^* = Np/(N - sp)$ is the fractional critical Sobolev exponent. Let us recall the weak formulation of problem (1.1). Let

$$[u]_{s,p} = \left(\int_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy \right)^{1/p}$$

be the Gagliardo seminorm of the measurable function $u : \mathbb{R}^N \rightarrow \mathbb{R}$, and let

$$W^{s,p}(\mathbb{R}^N) = \{u \in L^p(\mathbb{R}^N) : [u]_{s,p} < \infty\}$$

be the fractional Sobolev space endowed with the norm

$$\|u\|_{s,p} = (|u|_p^p + [u]_{s,p}^p)^{1/p},$$

where $|\cdot|_p$ is the norm in $L^p(\mathbb{R}^N)$ (see Di Nezza et al. [16] for details). We work in the closed linear subspace

$$X_p^s(\Omega) = \{u \in W^{s,p}(\mathbb{R}^N) : u = 0 \text{ a.e. in } \mathbb{R}^N \setminus \Omega\},$$

equivalently renormed by setting $\|\cdot\| = [\cdot]_{s,p}$, which is a uniformly convex Banach space. By [16, Theorems 6.5 & 7.1], the imbedding $X_p^s(\Omega) \hookrightarrow L^r(\Omega)$ is continuous for $r \in [1, p_s^*]$ and compact for $r \in [1, p_s^*)$. We let

$$S_{s,p} = \inf_{u \in X_p^s(\Omega) \setminus \{0\}} \frac{\|u\|^p}{|u|_{p_s^*}^p} \quad (1.2)$$

denote the best imbedding constant when $r = p_s^*$. A function $u \in X_p^s(\Omega)$ is a weak solution of problem (1.1) if

$$\begin{aligned} \int_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y)) (v(x) - v(y))}{|x - y|^{N+sp}} dx dy \\ = \lambda \int_{\Omega} |u|^{p-2} uv dx + \int_{\Omega} |u|^{p_s^*-2} uv dx \quad \forall v \in X_p^s(\Omega). \end{aligned} \quad (1.3)$$

In the semilinear case $p = 2$, problem (1.1) reduces to the critical fractional Laplacian problem

$$\begin{cases} (-\Delta)^s u = \lambda u + |u|^{2_s^*-2} u & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^N \setminus \Omega, \end{cases} \quad (1.4)$$

where $2_s^* = 2N/(N-2s)$. This nonlocal problem generalizes the well-known Brézis-Nirenberg problem, which has been extensively studied beginning with the seminal paper [8] (see, e.g., [11, 10, 1, 25, 21, 42, 12, 17, 22, 24, 23, 13, 40] and references therein). Consequently, many results known in the local case $s = 1$ have been extended to problem (1.4) (see, e.g., [38, 36, 39, 37, 2, 41, 19]). In particular, Fiscella et al. [19] have recently obtained the following bifurcation and multiplicity result, extending a well-known result of Cerami et al. [11] in the local case. Let $0 < \lambda_1 < \lambda_2 \leq \lambda_3 \leq \dots \rightarrow +\infty$ be the eigenvalues of the problem

$$\begin{cases} (-\Delta)^s u = \lambda u & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^N \setminus \Omega, \end{cases}$$

repeated according to multiplicity, and let $|\cdot|$ denote the Lebesgue measure in \mathbb{R}^N . If $\lambda_k \leq \lambda < \lambda_{k+1}$ and

$$\lambda > \lambda_{k+1} - \frac{S_{s,2}}{|\Omega|^{2s/N}},$$

and m denotes the multiplicity of λ_{k+1} , then problem (1.4) has m distinct pairs of nontrivial solutions $\pm u_j^\lambda$, $j = 1, \dots, m$ such that $u_j^\lambda \rightarrow 0$ as $\lambda \nearrow \lambda_{k+1}$ (see [19, Theorem 1]).

In the present paper we extend the above bifurcation and multiplicity result to the quasilinear nonlocal problem (1.1). This extension is quite nontrivial. Indeed, the linking argument based on eigenspaces of $(-\Delta)^s$ in [19] does not work when $p \neq 2$ since the nonlinear operator $(-\Delta)_p^s$ does not have linear eigenspaces. We will use a more general construction based on sublevel sets as in Perera and Szulkin [35] (see also Perera et al. [33, Proposition 3.23]). Moreover, the standard sequence of variational eigenvalues of $(-\Delta)_p^s$ based on the genus does not give enough information about the structure of the sublevel sets to carry out this linking construction. Therefore we will use a different sequence of eigenvalues introduced in Iannizzotto et al. [27] that is based on the \mathbb{Z}_2 -cohomological index of Fadell and Rabinowitz [18], which is defined as follows. Let W be a Banach space and let \mathcal{A} denote the class of symmetric subsets of $W \setminus \{0\}$. For $A \in \mathcal{A}$, let $\bar{A} = A/\mathbb{Z}_2$ be the quotient space of A with each u and $-u$ identified, let $f : \bar{A} \rightarrow \mathbb{R}P^\infty$ be the classifying map of \bar{A} , and let $f^* : H^*(\mathbb{R}P^\infty) \rightarrow H^*(\bar{A})$ be the induced homomorphism of the Alexander-Spanier cohomology rings. The cohomological index of A is defined by

$$i(A) = \begin{cases} \sup \{m \geq 1 : f^*(\omega^{m-1}) \neq 0\}, & A \neq \emptyset \\ 0, & A = \emptyset, \end{cases}$$

where $\omega \in H^1(\mathbb{R}P^\infty)$ is the generator of the polynomial ring $H^*(\mathbb{R}P^\infty) = \mathbb{Z}_2[\omega]$. For example, the classifying map of the unit sphere S^{m-1} in \mathbb{R}^m , $m \geq 1$ is the inclusion $\mathbb{R}P^{m-1} \subset \mathbb{R}P^\infty$, which induces isomorphisms on H^q for $q \leq m-1$, so $i(S^{m-1}) = m$.

The Dirichlet spectrum of $(-\Delta)_p^s$ in Ω consists of those $\lambda \in \mathbb{R}$ for which the problem

$$\begin{cases} (-\Delta)_p^s u = \lambda |u|^{p-2} u & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^N \setminus \Omega \end{cases} \quad (1.5)$$

has a nontrivial solution. Although a complete description of the spectrum is not known when $p \neq 2$, we can define an increasing and unbounded sequence of variational eigenvalues via a suitable minimax scheme. The standard scheme based on the genus does not give the index information necessary for our purposes here, so we will use the following scheme based on the cohomological index as in Iannizzotto et al. [27] (see also Perera [32]). Let

$$\Psi(u) = \frac{1}{|u|_p^p}, \quad u \in \mathcal{M} = \{u \in X_p^s(\Omega) : \|u\| = 1\}.$$

Then eigenvalues of problem (1.5) coincide with critical values of Ψ . We use the standard notation

$$\Psi^a = \{u \in \mathcal{M} : \Psi(u) \leq a\}, \quad \Psi_a = \{u \in \mathcal{M} : \Psi(u) \geq a\}, \quad a \in \mathbb{R}$$

for the sublevel sets and superlevel sets, respectively. Let \mathcal{F} denote the class of symmetric subsets of \mathcal{M} , and set

$$\lambda_k := \inf_{M \in \mathcal{F}, i(M) \geq k} \sup_{u \in M} \Psi(u), \quad k \in \mathbb{N}.$$

Then $0 < \lambda_1 < \lambda_2 \leq \lambda_3 \leq \dots \rightarrow +\infty$ is a sequence of eigenvalues of problem (1.5), and

$$\lambda_k < \lambda_{k+1} \implies i(\Psi^{\lambda_k}) = i(\mathcal{M} \setminus \Psi_{\lambda_{k+1}}) = k \quad (1.6)$$

(see Iannizzotto et al. [27, Proposition 2.4]). The asymptotic behavior of these eigenvalues was recently studied in Iannizzotto and Squassina [28]. Making essential use of the index information in (1.6), we will prove the following theorem.

Theorem 1.1. (i) *If*

$$\lambda_1 - \frac{S_{s,p}}{|\Omega|^{sp/N}} < \lambda < \lambda_1,$$

then problem (1.1) has a pair of nontrivial solutions $\pm u^\lambda$ such that $u^\lambda \rightarrow 0$ as $\lambda \nearrow \lambda_1$.

(ii) *If $\lambda_k \leq \lambda < \lambda_{k+1} = \dots = \lambda_{k+m} < \lambda_{k+m+1}$ for some $k, m \in \mathbb{N}$ and*

$$\lambda > \lambda_{k+1} - \frac{S_{s,p}}{|\Omega|^{sp/N}}, \quad (1.7)$$

then problem (1.1) has m distinct pairs of nontrivial solutions $\pm u_j^\lambda$, $j = 1, \dots, m$ such that $u_j^\lambda \rightarrow 0$ as $\lambda \nearrow \lambda_{k+1}$.

In particular, we have the following existence result.

Corollary 1.2. *Problem (1.1) has a nontrivial solution for all $\lambda \in \bigcup_{k=1}^{\infty} (\lambda_k - S_{s,p}/|\Omega|^{sp/N}, \lambda_k)$.*

We note that $\lambda_1 \geq S_{s,p}/|\Omega|^{sp/N}$. Indeed, if φ_1 is an eigenfunction associated with λ_1 ,

$$\lambda_1 = \frac{\|\varphi_1\|^p}{|\varphi_1|_p^p} \geq \frac{S_{s,p} |\varphi_1|_{p_s^*}^p}{|\varphi_1|_p^p} \geq \frac{S_{s,p}}{|\Omega|^{sp/N}}$$

by the Hölder inequality.

Remark 1.3. Analogous results for the corresponding local problem driven by the p -Laplacian operator were recently obtained in Perera et al. [34]. In this work new difficulties need to be handled due to the non-locality of the problem, in particular the detection of a (lower) range of validity of the Palais-Smale condition, consistent with the one known in the local case $s = 1$, see Proposition 3.1.

2 An abstract critical point theorem

The abstract result of Bartolo et al. [3] used in Cerami et al. [11] and Fiscella et al. [19] is based on linear subspaces and therefore cannot be used to prove our Theorem 1.1. In this section we prove a more general critical point theorem based on a pseudo-index related to the cohomological index that is applicable here (see also Perera et al. [33, Proposition 3.44]).

Let W be a Banach space and let \mathcal{A} denote the class of symmetric subsets of $W \setminus \{0\}$. The following proposition summarizes the basic properties of the cohomological index.

Proposition 2.1 (Fadell-Rabinowitz [18]). *The index $i : \mathcal{A} \rightarrow \mathbb{N} \cup \{0, \infty\}$ has the following properties:*

- (i₁) *Definiteness: $i(A) = 0$ if and only if $A = \emptyset$;*
- (i₂) *Monotonicity: If there is an odd continuous map from A to B (in particular, if $A \subset B$), then $i(A) \leq i(B)$. Thus, equality holds when the map is an odd homeomorphism;*
- (i₃) *Dimension: $i(A) \leq \dim W$;*
- (i₄) *Continuity: If A is closed, then there is a closed neighborhood $N \in \mathcal{A}$ of A such that $i(N) = i(A)$. When A is compact, N may be chosen to be a δ -neighborhood $N_\delta(A) = \{u \in W : \text{dist}(u, A) \leq \delta\}$;*
- (i₅) *Subadditivity: If A and B are closed, then $i(A \cup B) \leq i(A) + i(B)$;*
- (i₆) *Stability: If SA is the suspension of $A \neq \emptyset$, obtained as the quotient space of $A \times [-1, 1]$ with $A \times \{1\}$ and $A \times \{-1\}$ collapsed to different points, then $i(SA) = i(A) + 1$;*
- (i₇) *Piercing property: If A , A_0 and A_1 are closed, and $\varphi : A \times [0, 1] \rightarrow A_0 \cup A_1$ is a continuous map such that $\varphi(-u, t) = -\varphi(u, t)$ for all $(u, t) \in A \times [0, 1]$, $\varphi(A \times [0, 1])$ is closed, $\varphi(A \times \{0\}) \subset A_0$ and $\varphi(A \times \{1\}) \subset A_1$, then $i(\varphi(A \times [0, 1]) \cap A_0 \cap A_1) \geq i(A)$;*
- (i₈) *Neighborhood of zero: If U is a bounded closed symmetric neighborhood of 0, then $i(\partial U) = \dim W$.*

Let Φ be an even C^1 -functional defined on W , and recall that Φ satisfies the Palais-Smale compactness condition at the level $c \in \mathbb{R}$, or $(PS)_c$ for short, if every sequence $(u_j) \subset W$ such that $\Phi(u_j) \rightarrow c$ and $\Phi'(u_j) \rightarrow 0$ has a convergent subsequence. Let \mathcal{A}^* denote the class of symmetric subsets of W , let $r > 0$, let $S_r = \{u \in W : \|u\| = r\}$, let $0 < b \leq +\infty$, and let Γ denote the group of odd homeomorphisms of W that are the identity outside $\Phi^{-1}(0, b)$. The pseudo-index of $M \in \mathcal{A}^*$ related to i , S_r and Γ is defined by

$$i^*(M) = \min_{\gamma \in \Gamma} i(\gamma(M) \cap S_r)$$

(see Benci [4]). The following critical point theorem generalizes Bartolo et al. [3, Theorem 2.4].

Theorem 2.2. *Let A_0, B_0 be symmetric subsets of S_1 such that A_0 is compact, B_0 is closed, and*

$$i(A_0) \geq k + m, \quad i(S_1 \setminus B_0) \leq k$$

for some integers $k \geq 0$ and $m \geq 1$. Assume that there exists $R > r$ such that

$$\sup \Phi(A) \leq 0 < \inf \Phi(B), \quad \sup \Phi(X) < b,$$

where $A = \{Ru : u \in A_0\}$, $B = \{ru : u \in B_0\}$, and $X = \{tu : u \in A, 0 \leq t \leq 1\}$. For $j = k + 1, \dots, k + m$, let

$$\mathcal{A}_j^* = \{M \in \mathcal{A}^* : M \text{ is compact and } i^*(M) \geq j\}$$

and set

$$c_j^* := \inf_{M \in \mathcal{A}_j^*} \max_{u \in M} \Phi(u).$$

Then

$$\inf \Phi(B) \leq c_{k+1}^* \leq \dots \leq c_{k+m}^* \leq \sup \Phi(X),$$

in particular, $0 < c_j^ < b$. If, in addition, Φ satisfies the $(PS)_c$ condition for all $c \in (0, b)$, then each c_j^* is a critical value of Φ and there are m distinct pairs of associated critical points.*

Proof. If $M \in \mathcal{A}_{k+1}^*$,

$$i(S_r \setminus B) = i(S_1 \setminus B_0) \leq k < k + 1 \leq i^*(M) \leq i(M \cap S_r)$$

since $id_W \in \Gamma$. Hence M intersects B by (i_2) of Proposition 2.1. It follows that $c_{k+1}^* \geq \inf \Phi(B)$. If $\gamma \in \Gamma$, consider the continuous map

$$\varphi : A \times [0, 1] \rightarrow W, \quad \varphi(u, t) = \gamma(tu).$$

We have $\varphi(A \times [0, 1]) = \gamma(X)$, which is compact. Since γ is odd, $\varphi(-u, t) = -\varphi(u, t)$ for all $(u, t) \in A \times [0, 1]$ and $\varphi(A \times \{0\}) = \{\gamma(0)\} = \{0\}$. Since $\Phi \leq 0$ on A , $\gamma|_A = id_A$ and hence $\varphi(A \times \{1\}) = A$. Applying (i_7) with $\tilde{A}_0 = \{u \in W : \|u\| \leq r\}$ and $\tilde{A}_1 = \{u \in W : \|u\| \geq r\}$ gives

$$i(\gamma(X) \cap S_r) = i(\varphi(A \times [0, 1]) \cap \tilde{A}_0 \cap \tilde{A}_1) \geq i(A) = i(A_0) \geq k + m.$$

It follows that $i^*(X) \geq k + m$. So $X \in \mathcal{A}_{k+m}^*$ and hence $c_{k+m}^* \leq \sup \Phi(X)$. The rest now follows from standard arguments in critical point theory (see, e.g., Perera et al. [33]). \square

Remark 2.3. Constructions similar to the one in the proof of Theorem 2.2 have been used in Fadell and Rabinowitz [18] to prove bifurcation results for Hamiltonian systems, and in Perera and Szulkin [35] to obtain nontrivial solutions of p -Laplacian problems with nonlinearities that interact with the spectrum. See also Perera et al. [33, Proposition 3.44].

3 Proof of Theorem 1.1

Weak solutions of problem (1.1) coincide with critical points of the C^1 -functional

$$I_\lambda(u) = \frac{1}{p} \|u\|^p - \frac{\lambda}{p} |u|_p^p - \frac{1}{p_s^*} |u|_{p_s^*}^{p_s^*}, \quad u \in X_p^s(\Omega).$$

We have the following compactness result, which is well-known in the local case $s = 1$.

Proposition 3.1. *For any $\lambda \in \mathbb{R}$, I_λ satisfies the $(PS)_c$ condition for all $c < \frac{s}{N} S_{s,p}^{N/sp}$.*

First we prove a lemma.

Lemma 3.2. *If (u_j) is bounded in $X_p^s(\Omega)$ and $u_j \rightarrow u$ a.e. in Ω , then*

$$\|u_j\|^p = \|u_j - u\|^p + \|u\|^p + o(1) \quad \text{as } j \rightarrow \infty.$$

Proof. Defining $\omega_j : \mathbb{R}^{2N} \rightarrow \mathbb{R}_+$ by

$$\omega_j(x, y) = \left| \frac{|u_j(x) - u_j(y)|^p}{|x - y|^{N+sp}} - \frac{|(u_j(x) - u(x)) - (u_j(y) - u(y))|^p}{|x - y|^{N+sp}} - \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} \right|,$$

we will show that

$$\lim_{j \rightarrow \infty} \int_{\mathbb{R}^{2N}} \omega_j(x, y) \, dx dy = 0. \tag{3.1}$$

Given $\varepsilon > 0$, there exists $C_\varepsilon > 0$ such that

$$\left| |a + b|^p - |a|^p \right| \leq \varepsilon |a|^p + C_\varepsilon |b|^p \quad \forall a, b \in \mathbb{R},$$

and taking $a = (u_j(x) - u_j(y)) - (u(x) - u(y))$ and $b = u(x) - u(y)$ gives

$$\omega_j(x, y) \leq \varepsilon \frac{|(u_j(x) - u_j(y)) - (u(x) - u(y))|^p}{|x - y|^{N+sp}} + C_\varepsilon \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}}.$$

Consequently, defining $\omega_j^\varepsilon : \mathbb{R}^{2N} \rightarrow \mathbb{R}_+$ by

$$\omega_j^\varepsilon(x, y) = \left(\omega_j(x, y) - \varepsilon \frac{|(u_j(x) - u_j(y)) - (u(x) - u(y))|^p}{|x - y|^{N+sp}} \right)^+,$$

we have

$$\omega_j^\varepsilon(x, y) \leq C_\varepsilon \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} \in L^1(\mathbb{R}^{2N}).$$

Since $u_j \rightarrow u$ a.e. in \mathbb{R}^N , $\omega_j^\varepsilon \rightarrow 0$ a.e. in \mathbb{R}^{2N} , so the dominated convergence theorem now implies

$$\lim_{j \rightarrow \infty} \int_{\mathbb{R}^{2N}} \omega_j^\varepsilon(x, y) dx dy = 0.$$

Then

$$\limsup_{j \rightarrow \infty} \int_{\mathbb{R}^{2N}} \omega_j(x, y) dx dy \leq \varepsilon \limsup_{j \rightarrow \infty} \int_{\mathbb{R}^{2N}} \frac{|(u_j(x) - u_j(y)) - (u(x) - u(y))|^p}{|x - y|^{N+sp}} dx dy,$$

and (3.1) follows since $\varepsilon > 0$ is arbitrary and u_j is bounded in $X_p^s(\Omega)$. \square

Proof of Proposition 3.1. Let $c < \frac{s}{N} S_{s,p}^{N/sp}$ and let (u_j) be a sequence in $X_p^s(\Omega)$ such that

$$I_\lambda(u_j) = \frac{1}{p} \|u_j\|^p - \frac{\lambda}{p} |u_j|_p^p - \frac{1}{p_s^*} |u_j|_{p_s^*}^{p_s^*} = c + o(1), \quad (3.2)$$

$$\begin{aligned} I'_\lambda(u_j) v &= \int_{\mathbb{R}^{2N}} \frac{|u_j(x) - u_j(y)|^{p-2} (u_j(x) - u_j(y)) (v(x) - v(y))}{|x - y|^{N+sp}} dx dy \\ &\quad - \lambda \int_{\Omega} |u_j|^{p-2} u_j v dx - \int_{\Omega} |u_j|^{p_s^*-2} u_j v dx = o(\|v\|) \quad \forall v \in X_p^s(\Omega) \end{aligned} \quad (3.3)$$

as $j \rightarrow \infty$. Then

$$\frac{s}{N} |u_j|_{p_s^*}^{p_s^*} = I_\lambda(u_j) - \frac{1}{p} I'_\lambda(u_j) u_j = o(\|u_j\|) + \mathcal{O}(1),$$

which together with (3.2) and the Hölder inequality shows that (u_j) is bounded in $X_p^s(\Omega)$. So a renamed subsequence of (u_j) converges to some u weakly in $X_p^s(\Omega)$, strongly in $L^r(\Omega)$ for all $r \in [1, p_s^*)$, and a.e. in Ω (see Di Nezza et al. [16, Corollary 7.2]). Denoting by $p' = p/(p-1)$ the Hölder conjugate of p , $|u_j(x) - u_j(y)|^{p-2} (u_j(x) - u_j(y))/|x - y|^{(N+sp)/p'}$ is bounded in $L^{p'}(\mathbb{R}^{2N})$ and converges to $|u(x) - u(y)|^{p-2} (u(x) - u(y))/|x - y|^{(N+sp)/p'}$ a.e. in \mathbb{R}^{2N} , and $(v(x) - v(y))/|x - y|^{(N+sp)/p} \in L^p(\mathbb{R}^{2N})$, so the first integral in (3.3) converges to

$$\int_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y)) (v(x) - v(y))}{|x - y|^{N+sp}} dx dy$$

for a further subsequence. Moreover,

$$\int_{\Omega} |u_j|^{p-2} u_j v dx \rightarrow \int_{\Omega} |u|^{p-2} uv dx,$$

and

$$\int_{\Omega} |u_j|^{p_s^*-2} u_j v dx \rightarrow \int_{\Omega} |u|^{p_s^*-2} uv dx$$

since $|u_j|^{p_s^*-2} u_j \rightharpoonup |u|^{p_s^*-2} u$ in $L^{(p_s^*)'}(\Omega)$. So passing to the limit in (3.3) shows that $u \in X_p^s(\Omega)$ is a weak solution of (1.1), i.e., (1.3) holds.

Setting $\tilde{u}_j = u_j - u$, we will show that $\tilde{u}_j \rightarrow 0$ in $X_p^s(\Omega)$. We have

$$\|\tilde{u}_j\|^p = \|u_j\|^p - \|u\|^p + o(1) \quad (3.4)$$

by Lemma 3.2, and

$$|\tilde{u}_j|_{p_s^*}^{p_s^*} = |u_j|_{p_s^*}^{p_s^*} - |u|_{p_s^*}^{p_s^*} + o(1) \quad (3.5)$$

by the Brézis-Lieb lemma [7, Theorem 1]. Taking $v = u_j$ in (3.3) gives

$$\|u_j\|^p = \lambda |u|_p^p + |u_j|_{p_s^*}^{p_s^*} + o(1) \quad (3.6)$$

since (u_j) is bounded in $X_p^s(\Omega)$ and converges to u in $L^p(\Omega)$, and testing (1.3) with $v = u$ gives

$$\|u\|^p = \lambda |u|_p^p + |u|_{p_s^*}^{p_s^*}. \quad (3.7)$$

It follows from (3.4)–(3.7) and (1.2) that

$$\|\tilde{u}_j\|^p = |\tilde{u}_j|_{p_s^*}^{p_s^*} + o(1) \leq \frac{\|\tilde{u}_j\|_{p_s^*}^{p_s^*}}{S_{s,p}^{p_s^*/p}} + o(1),$$

so

$$\|\tilde{u}_j\|^p (S_{s,p}^{p_s^*/p} - \|\tilde{u}_j\|^{p_s^*-p}) \leq o(1). \quad (3.8)$$

On the other hand,

$$\begin{aligned} c &= \frac{1}{p} \|u_j\|^p - \frac{\lambda}{p} |u|_p^p - \frac{1}{p_s^*} \|u_j\|_{p_s^*}^{p_s^*} + o(1) \quad \text{by (3.2)} \\ &= \frac{s}{N} (\|u_j\|^p - \lambda |u|_p^p) + o(1) \quad \text{by (3.6)} \\ &= \frac{s}{N} (\|\tilde{u}_j\|^p + \|u\|^p - \lambda |u|_p^p) + o(1) \quad \text{by (3.4)} \\ &= \frac{s}{N} (\|\tilde{u}_j\|^p + |u|_{p_s^*}^{p_s^*}) + o(1) \quad \text{by (3.7)} \\ &\geq \frac{s}{N} \|\tilde{u}_j\|^p + o(1), \end{aligned}$$

so

$$\limsup_{j \rightarrow \infty} \|\tilde{u}_j\|^p \leq \frac{Nc}{s} < S_{s,p}^{N/sp}. \quad (3.9)$$

It follows from (3.8) and (3.9) that $\|\tilde{u}_j\| \rightarrow 0$. \square

If $\lambda_{k+m} < \lambda_{k+m+1}$, then $i(\Psi^{\lambda_{k+m}}) = k + m$ by (1.6). In order to apply Theorem 2.2 to the functional I_λ to prove Theorem 1.1, we will construct a compact symmetric subset A_0 of $\Psi^{\lambda_{k+m}}$ with the same index. As noted in Iannizzotto et al. [27], the operator $A_p^s \in C(X_p^s(\Omega), X_p^s(\Omega)^*)$, where $X_p^s(\Omega)^*$ is the dual of $X_p^s(\Omega)$, defined by

$$A_p^s(u) v = \int_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y)) (v(x) - v(y))}{|x - y|^{N+sp}} dx dy, \quad u, v \in X_p^s(\Omega)$$

satisfies the structural assumptions of [33, Chapter 1]. In particular, A_p^s is of type (S), i.e., every sequence $(u_j) \subset X_p^s(\Omega)$ such that

$$u_j \rightharpoonup u, \quad A_p^s(u_j) (u_j - u) \rightarrow 0$$

has a subsequence that converges strongly to u .

Lemma 3.3. *The operator A_p^s is strictly monotone, i.e.,*

$$(A_p^s(u) - A_p^s(v)) (u - v) > 0$$

for all $u \neq v$ in $X_p^s(\Omega)$.

Proof. By Perera et al. [33, Lemma 6.3], it suffices to show that

$$A_p^s(u) v \leq \|u\|^{p-1} \|v\| \quad \forall u, v \in X_p^s(\Omega)$$

and the equality holds if and only if $\alpha u = \beta v$ for some $\alpha, \beta \geq 0$, not both zero. We have

$$A_p^s(u) v \leq \int_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^{p-1} |v(x) - v(y)|}{|x - y|^{N+sp}} dx dy \leq \|u\|^{p-1} \|v\|$$

by the Hölder inequality. Clearly, equality holds throughout if $\alpha u = \beta v$ for some $\alpha, \beta \geq 0$, not both zero. Conversely, if $A_p^s(u) v = \|u\|^{p-1} \|v\|$, equality holds in both inequalities. The equality in the second inequality gives

$$\alpha |u(x) - u(y)| = \beta |v(x) - v(y)| \quad \text{a.e. in } \mathbb{R}^{2N}$$

for some $\alpha, \beta \geq 0$, not both zero, and then the equality in the first inequality gives

$$\alpha (u(x) - u(y)) = \beta (v(x) - v(y)) \quad \text{a.e. in } \mathbb{R}^{2N}.$$

Since u and v vanish a.e. in $\mathbb{R}^N \setminus \Omega$, it follows that $\alpha u = \beta v$ a.e. in Ω . \square

Lemma 3.4. *For each $w \in L^p(\Omega)$, the problem*

$$\begin{cases} (-\Delta)_p^s u = |w|^{p-2} w & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^N \setminus \Omega \end{cases} \quad (3.10)$$

has a unique weak solution $u \in X_p^s(\Omega)$. Moreover, the map $J : L^p(\Omega) \rightarrow X_p^s(\Omega)$, $w \mapsto u$ is continuous.

Proof. The existence follows from a standard minimization argument, and the uniqueness is immediate from the strict monotonicity of the operator A_p^s . Let $w_j \rightarrow w$ in $L^p(\Omega)$ and let $u_j = J(w_j)$, so

$$A_p^s(u_j) v = \int_{\Omega} |w_j|^{p-2} w_j v \, dx \quad \forall v \in X_p^s(\Omega). \quad (3.11)$$

Testing with $v = u_j$ gives

$$\|u_j\|^p = \int_{\Omega} |w_j|^{p-2} w_j u_j \, dx \leq |w_j|_p^{p-1} |u_j|_p$$

by the Hölder inequality, which together with the continuity of the imbedding $X_p^s(\Omega) \hookrightarrow L^p(\Omega)$ shows that (u_j) is bounded in $X_p^s(\Omega)$. So a renamed subsequence of (u_j) converges to some u weakly in $X_p^s(\Omega)$, strongly in $L^p(\Omega)$, and a.e. in Ω . An argument similar to that in the proof of Proposition 3.1 shows that u is a weak solution of (3.10), so $u = J(w)$. Testing (3.11) with $u_j - u$ gives

$$A_p^s(u_j) (u_j - u) = \int_{\Omega} |w_j|^{p-2} w_j (u_j - u) \, dx \rightarrow 0,$$

so $u_j \rightarrow u$ for a further subsequence as A_p^s is of type (S). \square

Proposition 3.5. *If $\lambda_l < \lambda_{l+1}$, then Ψ^{λ_l} has a compact symmetric subset A_0 with $i(A_0) = l$.*

Proof. Let

$$\pi_p(u) = \frac{u}{|u|_p}, \quad u \in X_p^s(\Omega) \setminus \{0\}$$

be the radial projection onto $\mathcal{M}_p = \{u \in X_p^s(\Omega) : |u|_p = 1\}$, and let

$$A = \pi_p(\Psi^{\lambda_l}) = \{w \in \mathcal{M}_p : \|w\|^p \leq \lambda_l\}.$$

Then $i(A) = i(\Psi^{\lambda_l}) = l$ by (i_2) of Proposition 2.1 and (1.6). For $w \in A$, let $u = J(w)$, where J is the map defined in Lemma 3.4, so

$$A_p^s(u) v = \int_{\Omega} |w|^{p-2} w v \, dx \quad \forall v \in X_p^s(\Omega).$$

Testing with $v = u, w$ and using the Hölder inequality gives

$$\|u\|^p \leq |w|_p^{p-1} |u|_p = |u|_p, \quad 1 = A_p^s(u) w \leq \|u\|^{p-1} \|w\|,$$

so

$$\|\pi_p(u)\| = \frac{\|u\|}{|u|_p} \leq \|w\|$$

and hence $\pi_p(u) \in A$. Let $\tilde{J} = \pi_p \circ J$ and let $\tilde{A} = \tilde{J}(A) \subset A$. Since the imbedding $X_p^s(\Omega) \hookrightarrow L^p(\Omega)$ is compact and \tilde{J} is an odd continuous map from $L^p(\Omega)$ to $X_p^s(\Omega)$, then \tilde{A} is a compact set and $i(\tilde{A}) = i(A) = l$. Let

$$\pi(u) = \frac{u}{\|u\|}, \quad u \in X_p^s(\Omega) \setminus \{0\}$$

be the radial projection onto \mathcal{M} and let $A_0 = \pi(\tilde{A})$. Then $A_0 \subset \Psi^{\lambda_l}$ is compact and $i(A_0) = i(\tilde{A}) = l$. \square

We are now ready to prove Theorem 1.1.

Proof of Theorem 1.1. We only give the proof of (ii). Proof of (i) is similar and simpler. By Proposition 3.1, I_λ satisfies the (PS) $_c$ condition for all $c < \frac{s}{N} S_{s,p}^{N/sp}$, so we apply Theorem 2.2 with $b = \frac{s}{N} S_{s,p}^{N/sp}$. By Proposition 3.5, $\Psi^{\lambda_{k+m}}$ has a compact symmetric subset A_0 with

$$i(A_0) = k + m.$$

We take $B_0 = \Psi_{\lambda_{k+1}}$, so that

$$i(S_1 \setminus B_0) = k$$

by (1.6). Let $R > r > 0$ and let A , B and X be as in Theorem 2.2. For $u \in B_0$,

$$I_\lambda(ru) \geq \frac{r^p}{p} \left(1 - \frac{\lambda}{\lambda_{k+1}}\right) - \frac{r^{p_s^*}}{p_s^* S_{s,p}^{p_s^*/p}}$$

by (1.2). Since $\lambda < \lambda_{k+1}$ and $p_s^* > p$, it follows that $\inf I_\lambda(B) > 0$ if r is sufficiently small. For $u \in A_0 \subset \Psi^{\lambda_{k+1}}$,

$$I_\lambda(Ru) \leq \frac{R^p}{p} \left(1 - \frac{\lambda}{\lambda_{k+1}}\right) - \frac{R^{p_s^*}}{p_s^* |\Omega|^{sp_s^*/N} \lambda_{k+1}^{p_s^*/p}}$$

by the Hölder inequality, so there exists $R > r$ such that $I_\lambda \leq 0$ on A . For $u \in X$,

$$\begin{aligned} I_\lambda(u) &\leq \frac{\lambda_{k+1} - \lambda}{p} \int_\Omega |u|^p dx - \frac{1}{p_s^* |\Omega|^{sp_s^*/N}} \left(\int_\Omega |u|^p dx \right)^{p_s^*/p} \\ &\leq \sup_{\rho \geq 0} \left[\frac{(\lambda_{k+1} - \lambda) \rho}{p} - \frac{\rho^{p_s^*/p}}{p_s^* |\Omega|^{sp_s^*/N}} \right] \\ &= \frac{s}{N} |\Omega| (\lambda_{k+1} - \lambda)^{N/sp}. \end{aligned}$$

So

$$\sup I_\lambda(X) \leq \frac{s}{N} |\Omega| (\lambda_{k+1} - \lambda)^{N/sp} < \frac{s}{N} S_{s,p}^{N/sp}$$

by (1.7). Theorem 2.2 now gives m distinct pairs of (nontrivial) critical points $\pm u_j^\lambda$, $j = 1, \dots, m$ of I_λ such that

$$0 < I_\lambda(u_j^\lambda) \leq \frac{s}{N} |\Omega| (\lambda_{k+1} - \lambda)^{N/sp} \rightarrow 0 \text{ as } \lambda \nearrow \lambda_{k+1}. \quad (3.12)$$

Then

$$|u_j^\lambda|_{p_s^*}^{p_s^*} = \frac{N}{s} \left[I_\lambda(u_j^\lambda) - \frac{1}{p} I'_\lambda(u_j^\lambda) u_j^\lambda \right] = \frac{N}{s} I_\lambda(u_j^\lambda) \rightarrow 0$$

and hence $u_j^\lambda \rightarrow 0$ in $L^p(\Omega)$ also by the Hölder inequality, so

$$\|u_j^\lambda\|^p = p I_\lambda(u_j^\lambda) + \lambda |u_j^\lambda|_p^p + \frac{p}{p_s^*} |u_j^\lambda|_{p_s^*}^{p_s^*} \rightarrow 0. \quad \square$$

References

- [1] A. Ambrosetti and M. Struwe, A note on the problem $-\Delta u = \lambda u + u|u|^{2^*-2}$ *Manuscripta Math.* **54** 373–379, (1986).
- [2] B. Barrios, E. Colorado, A. de Pablo, and U. Sánchez, On some critical problems for the fractional Laplacian operator. *J. Differential Equations* **252** 6133–6162, (2012).
- [3] P. Bartolo, V. Benci, and D. Fortunato, Abstract critical point theorems and applications to some nonlinear problems with strong resonance at infinity, *Nonlinear Anal.* **7** 981–1012, (1983).
- [4] V. Benci, On critical point theory for indefinite functionals in the presence of symmetries, *Trans. Amer. Math. Soc.* **274** 533–572, (1982).
- [5] L. Brasco, E. Parini, The second eigenvalue of the fractional p -Laplacian, preprint.
- [6] L. Brasco, E. Parini and M. Squassina, Stability of variational eigenvalues for the fractional p -Laplacian, *Discrete Cont. Dyn. Systems A*, to appear.
- [7] H. Brézis and E. Lieb, A relation between pointwise convergence of functions and convergence of functionals. *Proc. Amer. Math. Soc.* **88** 486–490, (1983).
- [8] H. Brézis and L. Nirenberg, Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents, *Comm. Pure Appl. Math.* **36** 437–477, (1983).

- [9] L. Caffarelli, Non-local diffusions, drifts and games. In *Nonlinear Partial Differential Equations*, **7** Abel Symposia, 37–52, (2012).
- [10] A. Capozzi, D. Fortunato, and G. Palmieri, An existence result for nonlinear elliptic problems involving critical Sobolev exponent, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **2** 463–470, (1985).
- [11] G. Cerami, D. Fortunato, and M. Struwe, Bifurcation and multiplicity results for nonlinear elliptic problems involving critical Sobolev exponents, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **1** 341–350, (1984).
- [12] D. G. Costa and E. A. Silva, A note on problems involving critical Sobolev exponents, *Differential Integral Equations* **8** 673–679, (1995).
- [13] E. A. de B. Silva and S. H. M. Soares. Quasilinear Dirichlet problems in \mathbb{R}^n with critical growth. *Nonlinear Anal.* **43** 1–20, (2001).
- [14] A. Di Castro, T. Kuusi, and G. Palatucci, Local behavior of fractional p -minimizers, preprint.
- [15] A. Di Castro, T. Kuusi, and G. Palatucci, Nonlocal Harnack inequalities, *J. Funct. Anal.* **267** 1807–1836, (2014).
- [16] E. Di Nezza, G. Palatucci, and E. Valdinoci, Hitchhiker’s guide to the fractional Sobolev spaces. *Bull. Sci. Math.* **136** 521–573, (2012).
- [17] P. Drábek and Y. X. Huang, Multiplicity of positive solutions for some quasilinear elliptic equation in \mathbb{R}^N with critical Sobolev exponent. *J. Differential Equations* **140** 106–132, (1997).
- [18] E. R. Fadell and P. H. Rabinowitz, Generalized cohomological index theories for Lie group actions with an application to bifurcation questions for Hamiltonian systems, *Invent. Math.* **45** 139–174, (1978).
- [19] A. Fiscella, G. M. Bisci, and R. Servadei, Bifurcation and multiplicity results for critical nonlocal fractional Laplacian problems, preprint.
- [20] G. Franzina and G. Palatucci, Fractional p -eigenvalues, *Riv. Mat. Univ. Parma* **5** 315–328, (2014).
- [21] J. G. Azorero and I. P. Alonso, Multiplicity of solutions for elliptic problems with critical exponent or with a nonsymmetric term, *Trans. Amer. Math. Soc.* **323** 877–895, (1991).
- [22] F. Gazzola and B. Ruf, Lower-order perturbations of critical growth nonlinearities in semilinear elliptic equations, *Adv. Differential Equations* **2** 555–572, (1997).

- [23] N. Ghoussoub and C. Yuan, Multiple solutions for quasi-linear PDEs involving the critical Sobolev and Hardy exponents. *Trans. Amer. Math. Soc.* **352** 5703–5743, (2000).
- [24] J. V. Gonçalves and C. O. Alves. Existence of positive solutions for m -Laplacian equations in \mathbb{R}^N involving critical Sobolev exponents. *Nonlinear Anal.* **32** 53–70, (1998).
- [25] M. Guedda and L. Véron, Quasilinear elliptic equations involving critical Sobolev exponents, *Nonlinear Anal.* **13** 879–902, (1989).
- [26] A. Iannizzotto, S. Mosconi, M. Squassina, Global Hölder regularity for the fractional p -Laplacian, preprint
- [27] A. Iannizzotto, S. Liu, K. Perera, and M. Squassina, Existence results for fractional p -Laplacian problems via Morse theory, *Adv. Calc. Var.*, to appear.
- [28] A. Iannizzotto and M. Squassina, Weyl-type laws for fractional p -eigenvalue problems. *Asympt. Anal.* **88** 233–245, (2014).
- [29] T. Kuusi, G. Mingione and Y. Sire Y., Nonlocal equations with measure data, *Comm. Math. Phys.*, to appear.
- [30] E. Lindgren, Hölder estimates for viscosity solutions of equations of fractional p -Laplace type, preprint
- [31] E. Lindgren and P. Lindqvist, Fractional eigenvalues, *Calc. Var. Partial Differential Equations* **49** 795–826, (2014).
- [32] K. Perera, Nontrivial critical groups in p -Laplacian problems via the Yang index, *Topol. Methods Nonlinear Anal.* **21** 301–309, (2003).
- [33] K. Perera, R. P. Agarwal, and D. O’Regan, Morse theoretic aspects of p -Laplacian type operators, volume 161 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2010.
- [34] K. Perera, M. Squassina, and Y. Yang, Bifurcation and multiplicity results for critical p -Laplacian problems, *Topol. Methods Nonlinear Anal.*, to appear.
- [35] K. Perera and A. Szulkin, p -Laplacian problems where the nonlinearity crosses an eigenvalue, *Discrete Contin. Dyn. Syst.* **13** 743–753, (2005).
- [36] R. Servadei, A critical fractional laplace equation in the resonant case, *Topol. Methods Nonlinear Anal.* **43** 251–267, (2014).
- [37] R. Servadei and E. Valdinoci, The Brezis-Nirenberg result for the fractional laplacian, *Trans. Amer. Math. Soc.* **367** (2015), 67–102.

- [38] R. Servadei, The Yamabe equation in a non-local setting. *Adv. Nonlinear Anal.* **2** 235–270, (2013).
- [39] R. Servadei and E. Valdinoci, A Brezis-Nirenberg result for non-local critical equations in low dimension, *Commun. Pure Appl. Anal.* **12** 2445–2464, (2013).
- [40] E. A. B. Silva and M. S. Xavier, Multiplicity of solutions for quasilinear elliptic problems involving critical Sobolev exponents, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **20** 341–358, (2003).
- [41] J. G. Tan, The Brezis-Nirenberg type problem involving the square root of the Laplacian, *Calc. Var. Partial Differential Equations* **42** 21–41, (2011).
- [42] Z. H. Wei and X. M. Wu, A multiplicity result for quasilinear elliptic equations involving critical Sobolev exponents, *Nonlinear Anal.* **18** 559–567, (1992).